

TABLE OF CONTENTS

Chapter	Page
I. INTRODUCTION	
Overview	1
Molecular Genetics Approaches Used to Investigate Populations	5
Studies Describing Genetic Variation in Populations of Brook Trout	7
Distribution and Status of Brook Trout in New Jersey	17
Study Rationale and Research Objective	27
II. MATERIALS AND METHODS	
Study Design	29
Sample Collection	33
DNA Extraction	34
Microsatellite DNA Amplification	35
Fragment Analysis	35
Data Analysis	37
III. RESULTS	
Genetic Diversity	41
Population Structure	48
Individual Assignment	53
IV. DISCUSSION	
Genetic Diversity within Populations	56
Genetic Diversity among Populations and Genetic Structure	60
Management and Conservation Implications	65
LITERATURE CITED	70
APPENDIX	77

LIST OF FIGURES

Figure	Page
1. Distribution of brook trout in North America (from MacCrimmon and Campbell 1969).	3
2. Distribution of wild (spawning) brook trout populations in New Jersey, as documented by stream surveys conducted by the New Jersey Division of Fish and Wildlife from 1968 through 2003 (from Hamilton and Barno 2005).	19
3. Examples of color variation in wild brook trout from New Jersey streams.	21
4. New Jersey's physiographic provinces and freshwaters having self-sustaining salmonid populations (trout production waters), as documented through NJDFW surveys conducted from 1968 through 2003 (Hamilton and Barno 2005).	22
5. Limits of glaciation in New Jersey and nearby New York.	23
6. Distribution and assessment of the status of wild brook trout in the eastern United States (left), with detail provided for New Jersey (right)(Hudy et al. 2005).	25
7. Map indicating the location of 22 sites in New Jersey where brook trout, <i>Salvelinus fontinalis</i> , were collected in 2000.	30
8. Collection of brook trout using a backpack electrofisher.	33
9. Cardiac puncture technique used to obtain blood samples.	33
10. DNA extraction success was confirmed electrophoretically and photo-documented.	34
11. Hierarchical gene diversity analysis (AMOVA) of 22 spawning populations of brook trout from New Jersey for 12 microsatellite DNA loci ($P < 0.010$).	47
12. Neighbor-joining phenogram depicting genetic distance using the chord distance of Cavelli-Sforza and Edwards (1967), for brook collected from 22 streams and 1 hatchery in New Jersey.	52

LIST OF TABLES

Table	Page
1. Location and trout stocking history information for 23 brook trout collection sites in New Jersey.	31
2. Three master mixes used to amplify 13 microsatellite loci in 23 brook trout collections from New Jersey.	36
3. Summary of genetic diversity statistics for 23 collections of brook trout (<i>Salvelinus fontinalis</i>) from New Jersey surveyed at 13 microsatellite loci.	42
4. Frequency of 21 unique alleles found in 238 brook trout collected from 22 streams and a hatchery in New Jersey, surveyed at 13 microsatellite loci.	45
5. Frequency of 29 unique alleles found in 238 brook trout collected from 4 drainages and a hatchery in New Jersey, surveyed at 13 microsatellite loci.	46
6. Matrix of F_{ST} values (below the diagonal) and R_{ST} values (above the diagonal) for all pairwise comparisons among 23 brook trout collections from New Jersey.	49
7. Genetic distance (chord distance, Cavalli-Sforza and Edwards 1967) among 23 collections of brook trout from New Jersey using 13 microsatellite loci.	50
8. Results of maximum likelihood assignment tests for 23 brook trout collections from New Jersey using multilocus genotypes derived from 12 microsatellite DNA markers.	54
9. Results of maximum likelihood assignment tests for 23 brook trout collections from four New Jersey drainages and a hatchery using multilocus genotypes derived from 12 microsatellite DNA markers.	55