

For patients ≥ 6 weeks of age and < 44 weeks of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>18 Weeks</i>	<i>32 Weeks</i>	<i>56 Weeks</i>	<i>150 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>10 Weeks</i>	<i>14 Weeks</i>	<i>24 Weeks</i>	<i>94 Weeks</i>
DTaP	X	X	X		X	X	
Hib	X	X	X		X		
IPV (Polio)	X	X	X		X		
PCV	X	X	X			X	
MMR						X	X
Varicella						X	X
Hepatitis A						X	X
Hepatitis B	X	X		X*			

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients 44 weeks of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>	<i>56 Weeks</i>	<i>150 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>24 Weeks</i>	<i>94 Weeks</i>
DTaP	X	X	X		X	X	
Hib	X	X	X				
IPV (Polio)	X	X	X		X		
PCV	X	X	X	X			
MMR						X	X
Varicella						X	X
Hepatitis A						X	X
Hepatitis B	X	X		X*			

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients ≥ 45 weeks of age and < 48 weeks of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>	<i>56 Weeks</i>	<i>150 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>24 Weeks</i>	<i>94 Weeks</i>
DTaP	X	X	X		X	X	
Hib	X	X	X				
IPV (Polio)	X	X	X		X		
PCV	X	X		X			
MMR						X	X
Varicella						X	X
Hepatitis A						X	X
Hepatitis B	X	X		X*			

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients ≥ 48 weeks of age and < 52 weeks of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>	<i>56 Weeks</i>	<i>150 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>16 Weeks</i>	<i>24 Weeks</i>	<i>94 Weeks</i>
DTaP	X	X	X		X	X	
Hib	X	X	X				
IPV (Polio)	X	X	X		X		
PCV	X	X		X			
MMR						X	X
Varicella						X	X
Hepatitis A						X	X
Hepatitis B	X	X		X*			

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients ≥ 12 months of age and < 14 months of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>	<i>56 Weeks</i>	<i>150 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>24 Weeks</i>	<i>94 Weeks</i>
DTaP	X	X	X		X	X	
Hib	X		X				
IPV (Polio)	X	X	X		X		
PCV	X		X				
MMR	X						X
Varicella	X						X
Hepatitis A	X				X		
Hepatitis B	X	X		X*			

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients ≥ 14 months of age and < 23 months of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>	<i>56 Weeks</i>	<i>150 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>24 Weeks</i>	<i>94 Weeks</i>
DTaP	X	X	X		X	X	
Hib	X						
IPV (Polio)	X	X	X		X		
PCV	X		X				
MMR	X						X
Varicella	X						X
Hepatitis A	X				X		
Hepatitis B	X	X		X*			

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients ≥ 23 months of age and < 3 years and 4 months of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>	<i>56 Weeks</i>	<i>105 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>24 Weeks</i>	<i>81 Weeks</i>
DTaP	X	X	X		X	X	
Hib	X						
IPV (Polio)	X	X	X		X		
PCV	X						
MMR	X						X
Varicella	X						X
Hepatitis A	X				X		
Hepatitis B	X	X		X*			

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients ≥3 years and 4 months of age and <3 years and 10 months of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>
DTaP	X	X	X		X
Hib	X				
IPV (Polio)	X	X	X		X
PCV	X				
MMR	X				X
Varicella	X				X
Hepatitis A	X				X
Hepatitis B	X	X		X*	

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients ≥ 3 years and 10 months of age and < 5 years of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>
DTaP	X	X	X		X
Hib	X				
IPV (Polio)	X	X	X		
PCV	X				
MMR	X		X		
Varicella	X		X		
Hepatitis A	X				X
Hepatitis B	X	X		X*	

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients ≥ 5 years of age and < 6 years and 4 months of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>
DTaP	X	X	X		X
IPV (Polio)	X	X	X		
MMR	X	X			
Varicella	X	X			
Hepatitis A	X				X
Hepatitis B	X	X		X*	

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients ≥ 6 years and 4 months of age and < 6 years and 10 months of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>
DTaP	X	X	X		
Td					X
IPV (Polio)	X	X	X		
MMR	X	X			
Varicella	X	X			
Hepatitis A	X				X
Hepatitis B	X	X		X*	

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients ≥ 6 years and 10 months of age and < 6 years and 11 months of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>
DTaP	X	X			
Td			X		X
IPV (Polio)	X	X	X		
MMR	X	X			
Varicella	X	X			
Hepatitis A	X				X
Hepatitis B	X	X		X*	

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients ≥ 6 years and 11 months of age and < 7 years of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>
DTaP	X				
Td		X	X		X
IPV (Polio)	X	X	X		
MMR	X	X			
Varicella	X	X			
Hepatitis A	X				X
Hepatitis B	X	X		X*	

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

For patients ≥ 7 years of age and < 9 years and 4 months of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>
Td	X	X			X
IPV (Polio)	X	X	X		
MMR	X	X			
Varicella	X	X			
Hepatitis A	X				X
Hepatitis B	X	X		X*	

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

Patients in this age group who do not receive a pertussis-containing vaccine during their initial Td series should receive a single Tdap vaccine at age 10 years.

For patients ≥ 9 years and 4 months of age and < 9 years and 11 months of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>
Td	X	X			
Tdap†					X
IPV (Polio)	X	X	X		
MMR	X	X			
Varicella	X	X			
Hepatitis A	X				X
Hepatitis B	X	X		X*	

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

† Boostrix® (licensed for patients 10 years of age) must be used for this Tdap vaccination

For patients ≥ 9 years 11 months of age and < 10 years of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>
Td	X				X
Tdap†		X			
IPV (Polio)	X	X	X		
MMR	X	X			
Varicella	X	X			
Hepatitis A	X				X
Hepatitis B	X	X		X*	

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

† Boostrix® (licensed for patients 10 years of age) must be used for this Tdap vaccination

For patients ≥ 10 years of age and < 18 years of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>
Td		X			X
Tdap†	X				
IPV (Polio)	X	X	X		
MMR	X	X			
Varicella	X	X			
Hepatitis A	X				X
Hepatitis B	X	X		X*	

* Third Hepatitis B vaccine must be given as a monovalent (noncombination) vaccine.

† Boostrix® (licensed for patients 10 years of age) must be used for this Tdap vaccination for patients < 11 years of age included on this table. Patients ≥ 11 years of age may receive either Boostrix® or Adacel®.

For patients ≥ 18 years of age AT THE INITIAL VISIT:

Time elapsed since initial vaccination	<i>Initial</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>	<i>32 Weeks</i>
Time from the last vaccine	<i>Visit</i>	<i>4 Weeks</i>	<i>4 Weeks</i>	<i>8 Weeks</i>	<i>16 Weeks</i>
Td		X			X
Tdap	X				
MMR	X	X			
Varicella	X	X			
Hepatitis A*	X	X			X
Hepatitis B*	X	X		X	

* If desired, combination Hepatitis A and B vaccine (Twinrix[®]) may be used in this age group. The combination vaccine should be administered on the same schedule as recommended for Hepatitis B alone (moving the 32-week Hepatitis A up to 16 weeks). If the combination vaccine is not used, the second Hepatitis A vaccination should be given at 32 weeks, and monovalent Hepatitis B vaccine should be used for the 16-week booster.