


Respiratory Virus Surveillance Report<sup>1</sup> New Jersey Department of Health Communicable Disease Service Week Ending January 21, 2017 (MMWR week 3)



# Synopsis




## ILI Activity<sup>4</sup>

		ILI Activity		
	Perce	nt ILI/Absenteeis	m	Baselines
	Current Week (range by county)	Last week Current year	Current week Last year	Non-season <sup>5</sup> Season <sup>6</sup> (3 low, 3 high)
Long Term Care Facilities	0.67 (0.00, 2.09)	0.63	0.68	0.56 (0.52, 0.76)
Schools (Absenteeism)	3.44 (1.29, 11.04)	5.53	5.69	3.42 (4.49, 4.81)
Emergency Departments	5.64 (0.96, 8.89)	4.75	3.49	2.35 (3.17, 3.81)
		Viral Activity	7	
		Current Week	Past 3 Weeks	Cumulative Total
Influenza H	H1N1 (2009)	0	0	1
Influen	za H3N2	31	157	299
Influ	enza B	4	32	65
	Syncytial Virus SV)	167	621	1813
Rapid Infl	uenza Tests	677	1815	2581


Report also available at http://nj.gov/health/flu/fluinfo.shtml

# Virologic Surveillance<sup>7</sup>

### Influenza Positive Specimens (PCR) – Result by Week




#### Influenza Positive Specimens (PCR) – Result by Region<sup>3</sup>




### Influenza Rapid Antigen Result by Week






# Influenza-Like Illness Surveillance



Respiratory Outbreaks in 1	Long Term Care Facilities <sup>9</sup>
Cumulative Outbreaks 2016-2017 season	34
No. outbreaks last 3 weeks	14
Regions with recent outbreaks	NE, CE, CW, S

### **ILINet Providers**



	ILINet F	Providers	
Curren	t Week	Previou	s Week
# of reporters	% ILI	# of reporters	% ILI
26	5.88	23	5.15

1	Pediatric Influenza Mortality <sup>1</sup>	0
Num	ber of Pediatric Influenza D Reported to CDC	leaths
Influenza Season	New Jersey	US (includes NJ)
2010 - 2011	4	123
2011 - 2012	1	35
2012 - 2013	7	171
2013 - 2014	6	108
2014 - 2015	1	146
2015 - 2016	1	85
2016 - 2017	0	5

For additional information regarding influenza surveillance please visit the following websites. <u>http://nj.gov/health/flu/surveillance.shtml</u> <u>http://www.cdc.gov/flu/</u>

### Footnotes:

- 1. This report represents activity occurring in New Jersey related to influenza and RSV. In addition, reports of other circulating respiratory viruses or regarding illness severity (i.e., hospitalization) will be included when available.
- 2. Activity levels for the state and region are defined in Table 1 and 2 at the end of this document.
- 3. The following is a breakdown of counties contained within each public health region: Northwest: Morris, Passaic, Sussex, Warren; Northeast: Bergen, Essex, Hudson; Central west: Hunterdon, Mercer, Somerset; Central East: Middlesex, Monmouth, Ocean, Union; South: Atlantic, Burlington, Camden, Cape May, Salem, Cumberland, Gloucester
- 4. Influenza-like illness (ILI) is defined as fever (> 100°F [37.8°C], oral or equivalent) and cough and/or sore throat (in the absence of a known cause other than influenza). For long term care facilities, fever is defined as 2°F above baseline temperature.
- 5. Non-season baseline is calculated by taking the average of statewide percentages of ILI for a 11 year (2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014,2015 and 2016) period during months when influenza is less likely to be circulating (May-August).
- 6. Three year seasonal averages are determined by calculating the average percent ILI/absenteeism for each influenza season (October to May). These averages are ranked and the three highest and lowest overall season averages were selected. The three highest and lowest numbers were then averaged to obtain a single high and single low value. The season which contribute to the high and low value vary by entity type and are as follows : LTCF (High: 10-11, 12-13, 14-15; Low: 11-12, 13-14, 15-16), ED (High: 12-13, 13-14, 14-15; Low: 10-11, 11-12, 15-16) and schools (High: 10-11, 12-13, 15-16; Low: 11-12,13-14, 14-15). A week by week average was also calculated using the average of the seasons listed above for each entity type.
- 7. Viral activity: Real-time polymerase chain reaction (PCR) results are obtained from electronic laboratory transmission submitted by acute care, commercial and public health laboratories to CDRSS. Rapid influenza test data and respiratory syncytial virus data are acquired from facilities reporting rapid influenza tests via the National Respiratory and Enteric Virus Surveillance System (NREVSS) or CDRSS ILI module. Counts for cumulative totals begin with week ending October 8, 2016. Three week count data includes current week and two prior weeks. Data presented for RSV and rapid influenza testing represent information for the week prior to the current report week.
- 8. Daily visits and admissions associated with ILI from emergency department data is collected via EpiCenter and Hippocrates. Prior to these systems, data on ILI visits were only recorded one day per week usually on Tuesday. This system is maintained as a large amount of historical data allows for better seasonal comparisons.
- 9. Only LTCF outbreaks reported to NJDOH that receive an outbreak number are recorded in this report.
- 10. Data presented for New Jersey are for cases confirmed as of the current reporting week. Data presented for the United States represent data reported for the prior MMWR week. This data can be viewed at: <u>http://www.cdc.gov/flu/weekly/</u>.

	Influenza Activity	<u>Table 1</u> Level – Definitions for State <b>2</b>	Activity	
NJ Level	CSTE Level		Definition	
		ILI Activity/Outbreaks		Lab Activity
	No Activity	ILI activity at or below baseline AND no detected outbreaks	AND	No lab confirmed cases
Low	Sporadic	Low ILI activity detected OR one lab confirmed outbreaks anywhere in the state	AND	Sporadic isolation of laboratory confirmed influenza
	Local	Increase in ILI activity OR two or more lab confirmed outbreaks in one public health region (Other regions not experiencing increased ILI activity)	AND	Recent (within 3 weeks) laboratory activity in the region with increased ILI
Moderate	Regional	Increase in ILI activity OR two or more lab confirmed outbreaks in at least 2 public health regions (Other regions not experiencing increased ILI activity)	AND	Recent (within 3 weeks) laboratory activity in the region with increased ILI
High	Widespread	Increase in ILI activity OR two or more lab confirmed outbreaks in > 2 public health regions	AND	Recent (within 3 weeks) laboratory activity in the region with increased ILI

	<u>Table 2</u> Influenza Activity Level – Definitions for Publ	ic Health	Regions
	<u> </u>	Definitio	<u>n</u>
<u>NJ Level</u>	ILI Activity/Outbreaks		Lab Activity
Low	Low ILI activity detected OR one lab confirmed outbreaks anywhere in the region	AND	Sporadic isolation of laboratory confirmed influenza anywhere in the region
Moderate	Increased ILI activity in less than half of the counties in the region OR two lab confirmed outbreaks in the public health region	AND	Recent (within 3 weeks) laboratory activity in same counties of the region with increased ILI
High	Increased ILI activity in more than half of the counties in the region OR three or more lab confirmed outbreaks in the region	AND	Recent (within 3 weeks) laboratory activity in more than half of the counties in the region with increased ILI

Notes:

ILI activity: Systems used to detect increases in ILI activity include: ILINet (i.e., sentinel providers), school absenteeism data, ED ILI visits and admissions collected via Hippocrates and EpiCenter systems, LTCF ILI data, LTCF outbreak data, and information on influenza mortality (122 city, influenza associated death report). Lab Activity: Virologic surveillance data from PHEL and commercial laboratories will be used as the primary data source for the above levels. However, rapid influenza test data will also be considered when determining the appropriate activity levels.

## INFLUENZA LABORATORY REPORTS BY COUNTY

# Counts represent total positive specimens from week ending October 7, 2016 to current MMWR week

## Source: CDRSS

Frequency			R	ESULT		
	COUNTY(COUNTY)	Influenza A - Typing not performed	Influenza A 2009 H1N1	Influenza AH3	Influenza B	Total
	ATLANTIC	104	0	9	5	118
	BERGEN	541	0	80	28	649
	BURLINGTON	59	0	4	9	72
	CAMDEN	252	0	12	8	272
	CAPE MAY	3	0	0	0	3
	CUMBERLAND	5	0	0	0	5
	ESSEX	96	0	16	13	125
	GLOUCESTER	75	0	2	4	81
	HUDSON	225	0	27	5	257
	HUNTERDON	35	0	7	0	42
	MERCER	66	0	4	2	72
	MIDDLESEX	75	0	13	4	92
	MONMOUTH	323	0	10	12	345
	MORRIS	119	0	35	7	161
	OCEAN	228	0	4	14	246
	PASSAIC	155	0	24	7	186
	SALEM	4	0	0	0	4
	SOMERSET	20	0	7	2	29
	SUSSEX	20	0	8	0	28
	UNION	118	1	42	15	176
	WARREN	24	0	3	2	29
	Total	2547	1	307	137	2992

## INFLUENZA LABORATORY REPORTS BY REGION

### Counts represent total positive specimens from week ending October 7, 2016 to current MMWR week

Frequency		Table of	REGION b	y RESULT		
			R	ESULT		
	REGION	Influenza A - Typing not performed	Influenza A 2009 H1N1	Influenza AH3	Influenza B	Total
	Central East	744	1	69	45	859
	Central West	121	0	18	4	143
	Northeast	862	0	123	46	1031
	Northwest	318	0	70	16	404
	South	502	0	27	26	555
	Total	2547	1	307	137	2992

### **Source: CDRSS**

The following is a breakdown of counties contained within each public health region: Northwest: Morris, Passaic, Sussex, Warren; Northeast: Bergen, Essex, Hudson Central west: Hunterdon, Mercer, Somerset Central East: Middlesex, Monmouth, Ocean, Union South: Atlantic, Burlington, Camden, Cape May, Salem, Cumberland, Gloucester


# NJ ACTIVE INFLUENZA-LIKE ILLNESS SURVEILLANCE STATISTICS SURVEILLANCE DATE: 01/17/2017





		Long Term Car	e		Schools		Hospi	tal Emergency	Dept
COUNTY	# Enrolled	# Reports Rec'd	% ILI	# Enrolled	# Reports Rec'd	% Absent	# Enrolled	# Reports Rec'd	% ILI
January 17, 2017 MMWR WEEK 3									
ATLANTIC	6	0	0.00	42	26	6.63	4	4	3.91
BERGEN	10	6	0.83	37	23	5.09	6	6	3.45
BURLINGTON	8	5	0.00	95	51	1.29	4	3	6.09
CAMDEN	1	0	0.00	1	0	0.00	7	7	4.70
CAPE MAY	3	0	0.00	14	8	5.61	1	1	5.69
CUMBERLAND	5	4	2.09	11	8	11.04	3	3	4.00
ESSEX	7	3	0.00	4	1	7.87	8	7	7.11
GLOUCESTER	3	1	0.00	4	1	3.92	2	2	2.69
HUDSON	4	1	0.87	14	7	8.17	6	6	8.13
HUNTERDON	4	3	1.08	8	7	5.08	1	1	3.49
MERCER	3	0	0.00	22	13	4.66	5	4	7.86
MIDDLESEX	11	5	0.00	22	17	4.98	6	6	5.52
MONMOUTH	6	2	0.00	16	5	6.05	5	5	5.59
MORRIS	2	0	0.00	9	5	2.61	4	4	3.76
OCEAN	8	4	1.06	5	3	7.27	4	4	5.25
PASSAIC	10	2	0.00	29	15	5.09	3	3	8.65
SALEM	0	0	0.00	3	3	6.87	1	1	7.35
SOMERSET	4	0	0.00	22	18	4.25	1	1	6.12
SUSSEX	2	1	0.00	5	3	7.94	2	2	0.96
UNION	1	0	0.00	51	17	4.70	5	5	4.29
WARREN	5	2	0.00	20	8	6.24	2	2	8.89
NW Region	19	5	0.00	63	31	5.16	11	11	6.36
NE Region	21	10	0.67	55	31	5.87	20	19	6.30
CW Region	11	3	1.08	52	38	4.54	7	6	7.09
CE Region	26	11	0.27	94	42	5.11	20	20	5.20
South Region	26	10	1.45	170	97	2.54	22	21	4.53
State Total	103	39	0.67	434	239	3.44	80	77	5.64

# NJ ACTIVE INFLUENZA-LIKE ILLNESS SURVEILLANCE STATISTICS SURVEILLANCE DATE: 01/17/2017





	RSV	' Tests	Rapid	Flu Tests
County	# Positive	Total Tests Performed	# Positive	Total Tests Performed
County January 17, 2017 MMWR WEEK		μ μ μ μ	#	μ μ μ μ
ATLANTIC	5	28	32	216
BERGEN	16	57	131	491
BURLINGTON	0	0	2	28
CAMDEN	0	6	35	428
CAMBEN CAPE MAY		10	1	37
	3 5	11	0	0
ESSEX	24	169	126	751
GLOUCESTER	7	32	32	287
HUDSON	5	34	45	161
HUNTERDON	2	9	45 12	88
MERCER	1	0	47	211
MIDDLESEX	3	49	10	81
MONMOUTH	35	192	139	819
MORRIS	33	192	14	102
OCEAN	2	18	35	208
PASSAIC	0	0	0	0
SALEM	0	0	0	0
SOMERSET	0	0	0	0
SUSSEX	13	77	16	83
UNION	13	114	0	24
WARREN	0	0	0	0
NW Region	46	275	30	185
NE Region	45	260	302	1403
_				
CW Region	3	9	59	299
CE Region	53	373	184	1132
South Region	20	87	102	996
State Total	167	1004	677	4015