


OHI #9: Hospitalizations from or with Pneumoconiosis

Annual Age-Standardized, Rate\* of Inpatient Hospitalizations from Other and Unspecified Pneumoconiosis, New Jersey and United States, 2000-2017



\*Age-Standardized rate per million residents

Estimated Annual Number and Age-Standardized Rate\* of Inpatient Hospitalizations from Other and Unspecified Pneumoconiosis, New Jersey and United States, 2000-2017

Year	Number		Rate*	
	NJ	US	NJ	US
2000	22	952	3.2	4.4
2001	28	709	4.1	3.2
2002	22	128	3.2	0.6
2003	23	418	3.2	1.9
2004	18	199	2.4	0.9
2005	12	1689	1.7	7.4
2006	17	1497	2.3	5.7
2007	22	1409	3.1	5.9
2008	25	N/A	3.6	N/A
2009	15	99	2.0	0.4
2010	9	1,220	1.2	4.7
2011	8	N/A	1.0	N/A
2012	7	N/A	1.0	N/A
2013	16	N/A	1.8	N/A
2014	12	N/A	1.6	N/A
2015	18	N/A	2.0	N/A
2016	17	N/A	2.0	N/A
2017	11	N/A	1.4	N/A

\*Age-standardized rate per million residents

N/A: not available

Note: Caution should be taken when comparing data to other states or with national estimates due to workers' compensation (WC) eligibility criteria and availability of data from WC programs varying among states. Effective October 1, 2015 healthcare organizations and providers were required to start using ICD-10-CM codes. The transition in classification from ICD-9-CM to ICD-10-CM may not be a one-to-one match, therefore caution should be taken when comparing data before and after the transition.

Data Source: NJ Department of Health inpatient hospital discharge data; US Census Bureau, American Fact Finder [[https://factfinder.census.gov/faces/nav/jsf/pages/index.xhtml?\\_ts=440355717799](https://factfinder.census.gov/faces/nav/jsf/pages/index.xhtml?_ts=440355717799)]; US 2000 Standard Population [<https://www.cdc.gov/nchs/data/statnt/statnt20.pdf>]