

**NJDEP**  
**Office of Science**  
**March 2014**

Preliminary Citation List for Interim Specific Ground Water Criterion for PFNA (March 13, 2014)

- Abbott, B.D., 2009. Review of the expression of peroxisome proliferator-activated receptors alpha (PPAR alpha), beta (PPAR beta), and gamma (PPAR gamma) in rodent and human development. *Reprod. Toxicol.* 27, 246–257.
- Abbott, B.D., Wood, C.R., Watkins, A.M., Das, K.P., Lau, C.S., 2010. Peroxisome proliferator-activated receptors alpha, beta, and gamma mRNA and protein expression in human fetal tissues. *PPAR Res.* pii: 690907, 26. Epub 2010 Jul.
- Barry V, Winquist A, Steenland K. Perfluorooctanoic acid (PFOA) exposures and incident cancers among adults living near a chemical plant. *Environ Health Perspect.* 2013 Nov-Dec;121(11-12):1313-8.
- Beesoon S, Webster GM, Shoeib M, Harner T, Benskin JP, Martin JW. Isomer profiles of perfluorochemicals in matched maternal, cord, and house dust samples: manufacturing sources and transplacental transfer. *Environ Health Perspect.* 2011 Nov;119(11):1659-64.
- Biegel, L.B., Hurtt, M.E., Frame, S.R., O'Connor, J.C., Cook, J.C., 2001. Mechanisms of extrahepatic tumor induction by peroxisome proliferators in male CD rats. *Toxicol. Sci.* 60, 44–55.
- Benninghoff, A.D., Bisson, W.H., Koch, D.C., Ehresman, D.J., Kolluri, S.K., Williams, D.E., 2011. Estrogen-like activity of perfluoroalkyl acids in vivo and interaction with human and rainbow trout estrogen receptors in vitro. *Toxicol. Sci.* 120, 42–58.
- Benninghoff AD, Orner GA, Buchner CH, Hendricks JD, Duffy AM, Williams DE. Promotion of hepatocarcinogenesis by perfluoroalkyl acids in rainbow trout. *Toxicol Sci.* 2012 Jan;125(1):69-78.
- C8 Science Panel website. C8 Science Panel reports and publications. <http://www.c8sciencepanel.org/>
- Das KP, Wood CR, Rosen MB, Lau C, and Abbott BD. 2013. Short-term exposure to perfluoroalkyl acids causes increase of hepatic lipid and triglyceride in conjunction with liver hypertrophy. *The Toxicologist*, Abstract #1981.
- Dong GH, Tung KY, Tsai CH, Liu MM, Wang D, Liu W, Jin YH, Hsieh WS, Lee YL, Chen PC. Serum polyfluoroalkyl concentrations, asthma outcomes, and immunological markers in a case-control study of Taiwanese children. *Environ Health Perspect.* 2013 Apr;121(4):507-13, 513e1-8.
- Emmett, E.A., Shofer, F.S., Zhang, H., Freeman, D., Desai, C., Shaw, L.M., 2006. Community exposure to perfluorooctanoate: Relationships between serum concentrations and exposure sources. *J. Occ. Environ. Med.* 48, 759–770.
- Fang X, Gao G, Xue H, Zhang X, Wang H. Exposure of perfluorononanoic acid suppresses the hepatic insulin signal pathway and increases serum glucose in rats. *Toxicology.* 2012 Apr 11;294(2-3):109-15.

Fang, X.; Zhang, L.; Feng, Y.; Zhao, Y.; Dai, J. Immunotoxic effects of perfluorononanoic acid on BALB/c mice. *Toxicol. Sci.* 2008, 105 (2), 312–321.

Feng Y, Fang X, Shi Z, Xu M, Dai J. Effects of PFNA exposure on expression of junction-associated molecules and secretory function in rat Sertoli cells. *Reprod Toxicol.* 2010 Nov;30(3):429-37.

Feng Y, Shi Z, Fang X, Xu M, Dai J. Perfluorononanoic acid induces apoptosis involving the Fas death receptor signaling pathway in rat testis. *Toxicol Lett.* 2009 Oct 28;190(2):224-30.

Fromme, H., Mosch, C., Morovitz, M., Alba-Alejandre, I., Boehmer, S., Kiranoglu, M., Faber, F., Hannibal, I., Genzel-Boroviczeny, O., Koletzko, B., Volk, W., 2010. Pre- and postnatal exposure to perfluorinated compounds (PFCs). *Environ. Sci. Technol.* 44, 7123–7129. and [Supporting Information](#).

Fujii Y, Yan J, Harada KH, Hitomi T, Yang H, Wang P, Koizumi A. Levels and profiles of long-chain perfluorinated carboxylic acids in human breast milk and infant formulas in East Asia. *Chemosphere.* 2012 Jan;86(3):315-21.

Granum B, Haug LS, Namork E, Stølevik SB, Thomsen C, Aaberge IS, van Loveren H, Løvik M, Nygaard UC. Pre-natal exposure to perfluoroalkyl substances may be associated with altered vaccine antibody levels and immune-related health outcomes in early childhood. *J Immunotoxicol.* 2013 Oct-Dec;10(4):373-9.

Gützkow KB, Haug LS, Thomsen C, Sabaredzovic A, Becher G, Brunborg G. Placental transfer of perfluorinated compounds is selective--a Norwegian Mother and Child sub-cohort study. *Int J Hyg Environ Health.* 2012 Feb;215(2):216-9.

Johansson, N., Fredriksson, A., Eriksson, P., 2008. Neonatal exposure to perfluorooctane sulfonate (PFOS) and perfluorooctanoic acid (PFOA) causes neurobehavioural defects in adult mice. *Neurotoxicology* 29, 160–169.

Johansson, N., Eriksson, P., Viberg, H., 2009. Neonatal exposure to PFOS and PFOA in mice results in changes in proteins which are important for neuronal growth and synaptogenesis in the developing brain. *Toxicol. Sci.* 108, 412–418.

Kato, K., Wong, L.Y., Jia, L.T., Kuklennyik, Z., Calafat, A.M., 2011. Trends in exposure to polyfluoroalkyl chemicals in the U.S. population: 1999–2008. *Environ. Sci. Technol.* 45, 8037–8045.

Kato, K., Calafat, A.M., Wong, L.Y., Wanigatunga, A.A., Caudill, S.P., Needham, L.L., 2009. Polyfluoroalkyl compounds in pooled sera from children participating in the National Health and Nutrition Examination Survey 2001–2002. *Environ. Sci. Technol.* 43, 2641–2647.

Kennedy GL Jr. Increase in mouse liver weight following feeding of ammonium perfluorooctanoate and related fluorochemicals. *Toxicol Lett.* 1987 Dec;39(2-3):295-300.

Kinney LA, Chromey NC, Kennedy GL Jr. Acute inhalation toxicity of ammonium perfluorononanoate. *Food Chem Toxicol.* 1989 Jul;27(7):465-8.

Kim, S.K., Lee, K.T., Kang, C.S., Tao, L., Kannan, K., Kim, K.R., Kim, C.K., Lee, J.S., Park, P.S., Yoo, Y.W., Ha, J.Y., Shin, Y.S., Lee, J.H., 2011. Distribution of perfluorochemicals between sera and milk from the same mothers and implications for prenatal and postnatal exposures. *Environ. Pollut.* 159, 169–174.

Lau, C.; Das, K. P.; Tatum, K.; Zehr, D.; Wood, C. R.; Rosen, M. B. Developmental toxicology of perfluorononanoic acid in the mouse. *The Toxicologist* 2009, 108, 417.

Lin CY, Wen LL, Lin LY, Wen TW, Lien GW, Hsu SH, Chien KL, Liao CC, Sung FC, Chen PC, Su TC. The associations between serum perfluorinated chemicals and thyroid function in adolescents and young adults. *J Hazard Mater.* 2013 Jan 15;244-245:637-44.

Lin CY, Wen LL, Lin LY, Wen TW, Lien GW, Chen CY, Hsu SH, Chien KL, Sung FC, Chen PC, Su TC. Associations between levels of serum perfluorinated chemicals and adiponectin in a young hypertension cohort in Taiwan. *Environ Sci Technol.* 2011 Dec 15;45(24):10691-8.

Lin CY, Chen PC, Lin YC, Lin LY. Association among serum perfluoroalkyl chemicals, glucose homeostasis, and metabolic syndrome in adolescents and adults. *Diabetes Care.* 2009 Apr;32(4):702-7.

Lind L, Zethelius B, Salihovic S, van Bavel B, Lind PM. Circulating levels of perfluoroalkyl substances and prevalent diabetes in the elderly. *Diabetologia.* 2014 Mar;57(3):473-9.

Liu, J., Li, J., Liu, Y., Chan, H.M., Zhao, Y., Cai, Z., Wu, Y., 2011. Comparison on gestation and lactation exposure of perfluorinated compounds for newborns. *Environ. Int.* 37, 1206–1212.

Lopez-Espinosa MJ, Mondal D, Armstrong B, Bloom MS, Fletcher T. Thyroid function and perfluoroalkyl acids in children living near a chemical plant. *Environ Health Perspect.* 2012 Jul;120(7):1036-41.

Macon, M.B., Villanueva, L.R., Tatum-Gibbs, K., Zehr, R.D., Strynar, M.J., Stanko, J.P., White, S.S., Helfant, L., Fenton, S.E., 2011. Prenatal perfluorooctanoic acid exposure in CD-1 mice: low dose developmental effects and internal dosimetry. *Toxicol. Sci.* 122, 134–145.

Mertens, J. J.; Sved, D. W.; Marit, G. B.; Myers, N. R.; Stetson, P. L.; Murphy, S. R.; Schmit, B.; Shinohara, M.; Farr, C. H. Subchronic toxicity of S-111-S-WB in Sprague Dawley rats. *Int. J. Toxicol.* 2010, 29 (4), 358–371.

Mondal D, Lopez-Espinosa MJ, Armstrong B, Stein CR, Fletcher T. Relationships of perfluorooctanoate and perfluorooctane sulfonate serum concentrations between mother-child pairs in a population with perfluorooctanoate exposure from drinking water. *Environ Health Perspect.* 2012 May;120(5):752-7.

Mondal D, Weldon RH, Armstrong BG, Gibson LJ, Lopez-Espinosa MJ, Shin HM, Fletcher T. Breastfeeding: a potential excretion route for mothers and implications for infant exposure to perfluoroalkyl acids. *Environ Health Perspect.* 2014 Feb;122(2):187-92.

Monroy R, Morrison K, Teo K, Atkinson S, Kubwabo C, Stewart B, Foster WG. Serum levels of perfluoroalkyl compounds in human maternal and umbilical cord blood samples. *Environ Res.* 2008 Sep;108(1):56-62.

Needham LL, Grandjean P, Heinzow B, Jørgensen PJ, Nielsen F, Patterson DG Jr, Sjödin A, Turner WE, Weihe P. Partition of environmental chemicals between maternal and fetal blood and tissues. *Environ Sci Technol.* 2011 Feb 1;45(3):1121-6.

Nelson JW, Hatch EE, Webster TF. Exposure to polyfluoroalkyl chemicals and cholesterol, body weight, and insulin resistance in the general U.S. population *Environ Health Perspect.* 2010 Feb;118(2):197-202.

NJDEP. 2007. Guidance for PFOA in Drinking Water at Pennsgrove Water Supply Company. [http://www.nj.gov/dep/watersupply/pdf/pfoa\\_dwguidance.pdf](http://www.nj.gov/dep/watersupply/pdf/pfoa_dwguidance.pdf)

Ode A, Rylander L, Lindh CH, Källén K, Jönsson BA, Gustafsson P, Olofsson P, Ivarsson SA, Rignell-Hydbom A. Determinants of maternal and fetal exposure and temporal trends of perfluorinated compounds. *Environ Sci Pollut Res Int*. 2013 Nov;20(11):7970-8.

Post, G.B., Louis, J.B., Lippincott, R.L., and Procopio, N.A. (2013). Occurrence of perfluorinated chemicals in raw water from New Jersey public drinking water systems. *Env. Sci. Technol.* 47(23):13266-75.

Post, G.B., Cohn, P.D., and Cooper, K.R. (2012). Perfluorooctanoic acid (PFOA), an emerging drinking water contaminant: a critical review of recent literature. *Env. Res.* 116: 93-117.

Post, G.B., Louis, J.B., Cooper, K.R., Boros-Russo, B.J., Lippincott, R.L., 2009. Occurrence and potential significance of perfluorooctanoic acid (PFOA) Detected in New Jersey public drinking water systems. *Environ. Sci. Technol.* 43, 4547–4554.

Post, G.B., Louis, J.B., Cooper, K.R., Lippincott, R.L., 2009. Response to comment on “Occurrence and Potential Significance of Perfluorooctanoic Acid (PFOA) Detected in New Jersey Public Drinking Water Systems.” *Environ. Sci. Technol.* 43, 8699–8700.

Rockwell CE, Turley AE, Cheng X, Fields PE, Klaassen CD (2013) Acute Immunotoxic Effects of Perfluorononanoic Acid (PFNA) in C57BL/6 Mice. *Clin Exp Pharmacol S4*: 002. doi:10.4172/2161-1459.S4-002

Rosen MB, Schmid JR, Zehr RD, Das KP, Ren H, Abbott BD, Lau C. 2010. Toxicogenomic profiling of perfluorononanoic acid in wild-type and PPAR $\alpha$ -null mice. *The Toxicologist*, Abstract 219.

Seals, R., Bartell, S.M., Steenland, K., 2011. Accumulation and clearance of perfluorooctanoic acid (PFOA) in current and former residents of an exposed community. *Environ. Health Perspect.* 119, 119–124.

Shin, H.M., Vieira, V.M., Ryan, P.B., Steenland, K., Bartell, S.M., 2011. Retrospective exposure estimation and predicted versus observed serum perfluorooctanoic acid concentrations for participants in the C8 Health Project. *Environ. Health Perspect.* 119, 1760–1765.

Sibinski, L.J., 1987. Final report of a two year oral (diet) toxicity and carcinogenicity study of fluorochemical FC-143 (perfluorooctanoate ammonium carboxylate) in rats. Vols. 1–4, 3M Company/RIKER. No.0281CR0012; 8EHQ-1087-0394, October 16, 1987.

Steenland, K., Jin, C., MacNeil, J., Lally, C., Ducatman, A., Vieira, V., Fletcher, T., 2009a. Predictors of PFOA levels in a community surrounding a chemical plant. *Environ. Health Perspect.* 117, 1083–1088.

Stump, D. G.; Holson, J. F.; Murphy, S. R.; Farr, C. H.; Schmit, B.; Shinohara, M. An oral two-generation reproductive toxicity study of S-111-S-WB in rats. *Reprod. Toxicol.* 2008, 25 (1), 7–20.

Takahashi M, Ishida S, Hirata-Koizumi M, Ono A, Hirose A. Repeated dose and reproductive/developmental toxicity of perfluoroundecanoic acid in rats. *J Toxicol Sci.* 2014;39(1):97-108.

Tao, L., Kannan, K., Wong, C.M., Arcaro, K.F., Butenhoff, J.L., 2008a. Perfluorinated compounds in human milk from Massachusetts, USA. *Environ. Sci. Technol.* 42, 3096–3101.

Tao, L., Ma, J., Kunisue, T., Libelo, E.L., Tanabe, S., Kannan, K., 2008b. Perfluorinated compounds in human breast milk from several Asian countries, and in infant formula and dairy milk from the United States. *Environ. Sci. Technol.* 42, 8597–8602.

Tatum-Gibbs, K.; Wambaugh, J. F.; Das, K. P.; Zehr, R. D.; Strynar, M. J.; Lindstrom, A. B.; Delinsky, A.; Lau, C. Comparative pharmacokinetics of perfluorononanoic acid in rat and mouse. *Toxicology* 2011, 281 (1–3), 48–55.

USEPA (2008). Child-Specific Exposure Factors Handbook. Chapter 3 -Water Ingestion. Table 3-1. Accessible at “Downloads” at <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=199243>

USEPA. 2009. United States Environmental Protection Agency. Existing Chemical Action Plans. Office of Pollution Prevention and Toxics. Long-Chain Perfluorinated Chemicals (PFCs) Action Plan. [http://www.epa.gov/opptintr/existingchemicals/pubs/pfcs\\_action\\_plan1230\\_09.pdf](http://www.epa.gov/opptintr/existingchemicals/pubs/pfcs_action_plan1230_09.pdf) 12/30/09

Vieira, V. M.; Hoffman, K.; Shin, H. M.; Weinberg, J. M.; Webster, T. F.; Fletcher, T. Perfluorooctanoic acid exposure and cancer outcomes in a contaminated community: A geographic analysis. *Environ. Health. Perspect.* 2013, 121 (3), 318–233.

White, S.S., Calafat, A.M., Kuklennyik, Z., Villanueva, L., Zehr, R.D., Helfant, L., Strynar, M.J., Lindstrom, A.B., Thibodeaux, J.R., Wood, C., Fenton, S.E., 2007. Gestational PFOA exposure of mice is associated with altered mammary gland development in dams and female offspring. *Toxicol. Sci.* 96, 133–144.

White, S.S., Kato, K., Jia, L.T., Basden, B.J., Calafat, A.M., Hines, E.P., Stanko, J.P., Wolf, C.J., Abbott, B.D., Fenton, S.E., 2009. Effects of perfluorooctanoic acid on mouse mammary gland development and differentiation resulting from cross-foster and restricted gestational exposures. *Reprod. Toxicol.* 27, 289–298.

White, S.S., Stanko, J.P., Kato, K., Calafat, A.M., Hines, E.P., Fenton, S.E., 2011b. Gestational and chronic low-dose PFOA exposures and mammary gland growth and differentiation in three generations of CD-1 Mice. *Environ. Health Perspect.* 119, 1070–1076.

Wolf, C. J.; Takacs, M. L.; Schmid, J. E.; Lau, C.; Abbott, B. D. Activation of mouse and human peroxisome proliferator-activated receptor alpha by perfluoroalkyl acids of different functional groups and chain lengths. *Toxicol. Sci.* 2008, 106 (1), 162–71.

Wolf, C.J., Schmid, J.E., Lau, C., Abbott, B.D. Activation of mouse and human peroxisome proliferator-activated receptor-alpha (PPAR $\alpha$ ) by perfluoroalkyl acids (PFAAs): Further investigation of C4–C12 compounds. *Reprod Toxicol.*, in press.

Wolf, C. J.; Zehr, R. D.; Schmid, J. E.; Lau, C.; Abbott, B. D. Developmental effects of perfluorononanoic acid in the mouse are dependent on peroxisome proliferator-activated receptor-alpha. *PPAR Res.* 2010, 2010, 282896.

Zhang T, Sun H, Lin Y, Qin X, Zhang Y, Geng X, Kannan K. Distribution of poly-and perfluoroalkyl substances in matched samples from pregnant women and carbon chain length related maternal transfer. *Environ Sci Technol.* 2013 Jul 16;47(14):7974-81.

Zhang Y, Beesoon S, Zhu L, Martin JW. Biomonitoring of perfluoroalkyl acids in human urine and estimates of biological half-life. *Environ Sci Technol*. 2013 Sep 17;47(18):10619-27.

DRAFT