


## Highlights

- **Influenza activity level is high statewide**
- **Emergency Department visits and outpatient provider visits associated with influenza-like illness are lower than last week & lower than this week last year**
- **There have been three confirmed influenza-associated pediatric deaths reported this season**
- **Outbreaks of Influenza continue to be reported in all settings throughout the state**
- **Positive results continue to be reported; influenza A (subtyping not performed) is at 73.16%, followed by A(H3N2) 22.71%, A (H1N1)pdm09 3.82%, and B 0.32%**


## Influenza Activity Level Comparison



## 1. Current Influenza Activity Level

This report summarizes surveillance information for influenza and other viral respiratory illnesses reported to the New Jersey Department of Health (NJDOH) Communicable Disease Service. As per regulation, influenza is a laboratory reportable condition but it is not possible to count every case that occurs since some individuals will not seek medical care or may never get tested. Surveillance is conducted year round and this report is published from October to May. The [Morbidity and Mortality Weekly Report \(MMWR\) week](#) is the time frame used by the Centers for Disease Control and Prevention (CDC) for disease reporting and activity levels are defined in the table on page 7 of this report. **Counts displayed below are the cumulative totals reported for the season beginning with MMWR week 40, week ending October 8, 2022.**

State Activity Level	
HIGH	
Regional Data	
<b>Northwest</b> Morris, Passaic, Sussex, Warren	HIGH
<b>Northeast</b> Bergen, Essex, Hudson	HIGH
<b>Central West</b> Hunterdon, Mercer, Somerset	HIGH
<b>Central East</b> Middlesex, Monmouth, Ocean, Union	HIGH
<b>Southwest</b> Burlington, Camden, Gloucester, Salem	HIGH
<b>Southeast</b> Atlantic, Cape May, Cumberland	HIGH



**75,786**  
Cases reported (PCR & Rapid)

**50**  
Outbreaks (Long Term Care)





**3**  
Pediatric flu deaths (confirmed)

## 2. Laboratory Testing

Real-time polymerase chain reaction (PCR) results for influenza (AH1N1, AH3N2, A subtyping not performed, and B) are obtained from electronic laboratory transmission submitted by acute care, commercial and public health laboratories. Rapid influenza test data are acquired from facilities reporting via the CDRSS Surveillance for Infectious Conditions (SIC) module. While the cumulative totals begin with MMWR week 40, week ending October 8, 2022, the data represented in charts begin with MMWR week 35, week ending September 3, 2022. Past 3 weeks data includes the current week and two prior weeks starting with MMWR week 40, week ending October 8, 2022.


	Test Type	Current Week	Past 3 Weeks	Cumulative Total
PCR	Influenza A (H1N1)pdm09	268	939	2498 (3.82%)
	Influenza A H3N2	494	2393	14856 (22.71%)
	Influenza A (Subtyping Not Performed)	3393	12815	47856 (73.16%)
	Influenza B	35	76	207 (0.32%)
Rapid	Rapid Influenza	770	3540	10369

## 3. Virologic Surveillance



### 3. Virologic Surveillance, *continued*

Data presented for rapid influenza testing represents information for the week prior to the current report week. Three year seasonal averages for rapid influenza tests are determined by calculating the average percent positive for each influenza season (October to May) beginning with the 2012-2013 season. These averages were ranked and the three highest and lowest overall season averages were selected. The three highest and lowest numbers were then averaged to obtain a single high and single low value for each week. The seasons which contribute to the high and low value for the rapid influenza chart are as follows: High: 16-17, 17-18, 18 -19; Low: 12-13, 14-15, 21-22. Off season baseline is calculated by taking the average of percent positivity for a 10 year period (2012 through and including 2022) during the months when influenza is less likely to be circulating (May to September). Data from the 19-20 and 20-21 seasons were excluded due to the COVID-19 pandemic.




### 4. Pediatric Influenza Mortality

Influenza-associated pediatric mortality was added to New Jersey’s reportable disease list in 2009. The below table includes severe and fatal influenza associated pediatric cases reported to NJDOH. Severe illness is defined as admission to an intensive care unit for an influenza-related illness. An influenza associated pediatric death is defined as a death resulting from a clinically compatible illness with lab confirmed influenza.

Influenza Season	US (fatal)	NJ (severe)	NJ (fatal)
2018-2019	106	51	6
2019-2020	188	57	2
2020-2021	1	1	0
2021-2022	43	19	0
2022-2023	61	44	3

### 5. Percent of Deaths due to Pneumonia and Influenza

Records of all deaths in New Jersey are maintained by NJDOH, Office of Vital Statistics and Registry and are submitted to the National Center for Health Statistics (NCHS). Pneumonia and influenza (P&I) deaths are identified from these records, compiled by the week of death and percentages are calculated. There is a 2-4 week lag period between the week the deaths have occurred and when the data for that week is reported. Because many influenza and COVID-19 deaths have pneumonia included on the death certificate, P&I no longer measures the impact of influenza in the same way as in the past. Additional information is also available at <https://gis.cdc.gov/grasp/fluview/mortality.html>.




## 6. ILI Activity

Influenza-like illness (ILI) is defined as fever (> 100°F [37.8°C], oral or equivalent) and cough and/or sore throat. For Long-term Care Facilities (LTCFs), fever is defined as 2°F above baseline temperature. ILI Activity from LTCFs and absenteeism data from schools is collected in the SIC Module of the Communicable Disease Reporting and Surveillance System (CDRSS). LTCFs and schools report their total census and number ill with ILI or number absent, respectively. Emergency department (ED) data is the aggregate weekly total of syndromic ILI visits and total ED registrations as recorded in EpiCenter (e.g., NJDOH syndromic surveillance system). Off season baseline is calculated by taking the average of statewide percentages of ILI for a 10 year period (2012 through and including 2022) during months when influenza is less likely to be circulating (May to September). Data from the 19-20 and 20-21 seasons were excluded due to the COVID-19 pandemic.

Percent Influenza-like Illness/Absenteeism				Baselines
	Current Week (range by county)	Last week Current year	Current week Last year	Off Season (Seasonal Average— low, high)
<b>Long-term Care Facilities</b>	0.15 (0.00, 1.18)	0.91	0.00	0.42 (0.30, 0.75)
<b>Emergency Departments</b>	6.77 (3.64, 14.78)	6.91	9.91	2.09 (3.52, 4.30)
<b>Schools (Absenteeism)</b>	CLOSED	6.22	CLOSED	3.98 (4.33, 4.93)

### 6a. Long-term Care Facility ILI Activity






### 6b. Long-term Care Facility Outbreaks

Only LTCF respiratory virus outbreaks reported to NJDOH that receive an outbreak number are recorded in this report. This does not include outbreaks due to COVID-19.


Respiratory Outbreaks in Long-term Care Facilities	
Cumulative Outbreaks 2022-2023 Season	50
No. outbreaks last 3 weeks	15
Regions with recent outbreaks	NW, NE, CW, CE, SW, SE

## 6c. Emergency Department ILI Activity (Syndromic Surveillance)

Daily visits and admissions associated with ILI from emergency department data are collected via EpiCenter (NJDOH syndromic surveillance). Prior to the 2017-2018 season, data on ILI visits were only recorded on one day per week usually on Tuesday. Beginning in the 2017-2018 season, weekly aggregate data is being recorded for ILI visits and admissions. Three year seasonal averages for emergency department visits and admissions are determined by calculating the average percent positivity for each influenza season (October to May) beginning with the 2012-2013 season. These averages were ranked and the three highest and lowest overall season averages were selected. The three highest and lowest numbers were then averaged to obtain a single high and single low value for each week. The seasons which contribute to the high and low value for emergency department visits chart are as follows: High: 12-13, 17-18, 18-19; Low: 13-14, 14-15, 15-16. The seasons which contribute to the high and low value for emergency department admissions chart are as follows: High: 13-14, 14-15, 17-18; Low: 12-13, 16-17, 21-22. Data from the 19-20 and 20-21 seasons were excluded due to the COVID-19 pandemic. Syndromic surveillance may capture other respiratory pathogens, such as SARS-CoV-2, that present with similar symptoms.




## 6d. School Absenteeism




## 7. ILI Net Providers

The U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) monitors outpatient visits for influenza-like illness (ILI), not laboratory-confirmed influenza, and may capture visits due to other respiratory pathogens, such as SARS-CoV-2, that present with similar symptoms.







## 8. Non-Influenza Viral Respiratory Surveillance

The National Respiratory and Enteric Virus Surveillance System (NREVSS) is a laboratory-based surveillance system and participating laboratories report the total number of tests performed and the total positive for a number of non-influenza respiratory viruses. Information about the CDC NREVSS system can be found at: <https://www.cdc.gov/surveillance/nrevss/labs/index.html>. Respiratory syncytial virus (RSV) data are acquired from facilities reporting via NREVSS or CDRSS SIC module. The RSV season is based upon the 5-year average of percent positivity and runs from the two consecutive weeks where percent positivity is at or above 10% through two consecutive weeks where it is below 10%.



## 8. Non-Influenza Viral Respiratory Surveillance (continued)



## Influenza Activity Level—Definitions for Public Health Regions

NJ Level	Definition		
	ILI Activity/Outbreaks		Lab Activity
<b>Low</b>	Low ILI activity detected OR one lab confirmed outbreak anywhere in the region	AND	Sporadic isolation of laboratory confirmed influenza anywhere in the region
<b>Moderate</b>	Increased ILI activity in less than half of the counties in the region OR two lab confirmed outbreaks in the public health region	AND	Recent (within 3 weeks) laboratory activity in the same counties of the region with increased ILI
<b>High</b>	Increased ILI activity in more than half of the counties in the region OR $\geq 3$ lab confirmed outbreaks in the region	AND	Recent (within 3 weeks) laboratory activity in more than half of the counties in the region with increased ILI

For additional information, visit the following websites: <http://nj.gov/health/flu/surveillance/shtml> and <http://www.cdc.gov/flu/>

## SURVEILLANCE DATE: 12/27/2022



COUNTY	Long Term Care			Schools			Hospital Emergency Dept		
	# Enrolled	# Reports Rec'd	% ILI	# Enrolled	# Reports Rec'd	% Absent	# Enrolled	# Reports Rec'd	% ILI
December 27, 2022 12:00 AM MMWR WEEK 52									
ATLANTIC	2	0	0.00	136	5	0.00	4	4	3.64
BERGEN	13	1	0.00	473	71	5.68	6	6	5.59
BURLINGTON	6	0	0.00	266	14	5.76	4	4	6.97
CAMDEN	1	0	0.00	239	50	0.27	7	7	6.99
CAPE MAY	3	0	0.00	50	0	0.00	1	1	5.41
CUMBERLAND	5	0	0.00	69	7	2.05	3	3	5.84
ESSEX	9	1	0.00	357	24	0.49	8	7	7.45
GLOUCESTER	3	0	0.00	117	2	0.00	3	3	8.50
HUDSON	4	0	0.00	259	66	0.94	6	6	8.57
HUNTERDON	4	0	0.00	68	10	0.10	1	1	4.72
MERCER	1	0	0.00	194	30	0.67	4	3	5.92
MIDDLESEX	14	0	0.00	337	43	0.32	6	6	8.08
MONMOUTH	6	0	0.00	342	41	12.33	5	5	5.70
MORRIS	3	0	0.00	239	25	2.30	4	4	5.89
OCEAN	9	1	0.00	316	9	0.85	4	4	4.32
PASSAIC	9	0	0.00	251	13	0.24	3	3	7.55
SALEM	0	0	0.00	40	3	0.00	1	1	14.78
SOMERSET	5	0	0.00	165	28	0.02	1	1	5.88
SUSSEX	3	0	0.00	63	4	0.00	1	1	5.87
UNION	3	0	0.00	313	20	0.32	5	5	8.47
WARREN	6	1	1.18	65	6	2.22	2	2	7.78
NW Region	21	1	1.18	618	48	1.32	10	10	10.77
NE Region	26	2	0.00	1089	161	2.28	20	19	7.11
CW Region	10	0	0.00	427	68	0.28	6	5	5.74
CE Region	32	1	0.00	1308	113	4.23	20	20	6.81
SW Region	10	0	0.00	662	69	1.90	15	15	9.62
SE Region	10	0	0.00	255	12	0.52	8	8	4.68
<b>State Total</b>	<b>109</b>	<b>4</b>	<b>0.15</b>	<b>4359</b>	<b>471</b>	<b>2.28</b>	<b>79</b>	<b>77</b>	<b>6.77</b>



## SURVEILLANCE DATE: 12/27/2022



County	RSV Tests		Rapid Flu Tests	
	# Positive	Total Tests Performed	# Positive	Total Tests Performed
December 27, 2022 12:00 AM MMWR WEEK 52				
ATLANTIC	11	909	134	1168
BERGEN	37	1165	81	500
BURLINGTON	0	0	14	77
CAMDEN	10	115	0	0
CAPE MAY	0	0	0	0
CUMBERLAND	8	1098	0	0
ESSEX	24	341	113	1341
GLOUCESTER	0	0	0	0
HUDSON	5	14	38	117
HUNTERDON	23	564	47	564
MERCER	0	0	0	0
MIDDLESEX	6	219	35	219
MONMOUTH	41	1352	279	2506
MORRIS	35	979	0	0
OCEAN	4	13	29	354
PASSAIC	20	300	0	0
SALEM	0	0	0	0
SOMERSET	326	4836	0	0
SUSSEX	0	0	0	0
UNION	22	1067	0	0
WARREN	0	0	0	0
NW Region	55	1279	0	0
NE Region	66	1520	232	1958
CW Region	349	5400	47	564
CE Region	73	2651	343	3079
SW Region	10	115	14	77
SE Region	19	2007	134	1168
<b>State Total</b>	<b>572</b>	<b>12972</b>	<b>770</b>	<b>6846</b>