

Respiratory Virus Surveillance Report<sup>1</sup> New Jersey Department of Health Communicable Disease Service Week Ending February 25, 2017 (MMWR week 8)



# Synopsis



#### **ILI Activity<sup>4</sup>**

		Ŭ		
	Perce	ent ILI/Absenteeis	m	Baselines
	Current Week (range by county)	Last week Current year	Current week Last year	Non-season <sup>5</sup> Season <sup>6</sup> (3 low, 3 high)
Long Term Care Facilities	0.43 (0.00, 4.45)	0.67	0.61	0.56 (0.52, 0.76)
Schools (Absenteeism)	5.77 (3.21, 9.38)	4.29	4.33	3.42 (4.49, 4.81)
Emergency Departments	5.20 (1.89, 8.60)	5.41	4.15	2.35 (3.17, 3.81)
		Viral Activity	7	
		Current Week	Past 3 Weeks	Cumulative Total
Influenza I	H1N1 (2009)	0	1	3
Influen	za H3N2	32	245	901
Influe	enza B	50	147	282
Respiratory S (R	byncytial Virus SV)	88	316	2562
Rapid Infl	uenza Tests	962	3350	8085

Report also available at http://nj.gov/health/flu/fluinfo.shtml

## Virologic Surveillance<sup>7</sup>

#### Influenza Positive Specimens (PCR) – Result by Week



#### Influenza Positive Specimens (PCR) – Result by Region<sup>3</sup>



#### Influenza Rapid Antigen Result by Week



#### Repiratory Syncytial Virus (RSV) Result by Week



# **Influenza-Like Illness Surveillance**



Respiratory Outbreaks in l	Long Term Care Facilities <sup>9</sup>
Cumulative Outbreaks 2016-2017 season	70
No. outbreaks last 3 weeks	14
Regions with recent outbreaks	NW, NE, CW, CE, S

#### **ILINet Providers**



	ILINet F	Providers	
Curren	t Week	Previou	ıs Week
# of reporters	% ILI	# of reporters	% ILI
25	4.84	29	5.59

]	Pediatric Influenza Mortality <sup>1</sup>	0
Numl	ber of Pediatric Influenza D Reported to CDC	leaths
Influenza Season	New Jersey	US (includes NJ)
2010 - 2011	4	123
2011 - 2012	1	35
2012 - 2013	7	171
2013 - 2014	6	108
2014 - 2015	1	146
2015 - 2016	1	85
2016 - 2017	0	34

For additional information regarding influenza surveillance please visit the following websites. <u>http://nj.gov/health/flu/surveillance.shtml</u> <u>http://www.cdc.gov/flu/</u>

#### Footnotes:

- 1. This report represents activity occurring in New Jersey related to influenza and RSV. In addition, reports of other circulating respiratory viruses or regarding illness severity (i.e., hospitalization) will be included when available.
- 2. Activity levels for the state and region are defined in Table 1 and 2 at the end of this document.
- 3. The following is a breakdown of counties contained within each public health region: Northwest: Morris, Passaic, Sussex, Warren; Northeast: Bergen, Essex, Hudson; Central west: Hunterdon, Mercer, Somerset; Central East: Middlesex, Monmouth, Ocean, Union; South: Atlantic, Burlington, Camden, Cape May, Salem, Cumberland, Gloucester
- 4. Influenza-like illness (ILI) is defined as fever (> 100°F [37.8°C], oral or equivalent) and cough and/or sore throat (in the absence of a known cause other than influenza). For long term care facilities, fever is defined as 2°F above baseline temperature.
- 5. Non-season baseline is calculated by taking the average of statewide percentages of ILI for a 11 year (2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014,2015 and 2016) period during months when influenza is less likely to be circulating (May-August).
- 6. Three year seasonal averages are determined by calculating the average percent ILI/absenteeism for each influenza season (October to May). These averages are ranked and the three highest and lowest overall season averages were selected. The three highest and lowest numbers were then averaged to obtain a single high and single low value. The season which contribute to the high and low value vary by entity type and are as follows : LTCF (High: 10-11, 12-13, 14-15; Low: 11-12, 13-14, 15-16), ED (High: 12-13, 13-14, 14-15; Low: 10-11, 11-12, 15-16) and schools (High: 10-11, 12-13, 15-16; Low: 11-12,13-14, 14-15). A week by week average was also calculated using the average of the seasons listed above for each entity type.
- 7. Viral activity: Real-time polymerase chain reaction (PCR) results are obtained from electronic laboratory transmission submitted by acute care, commercial and public health laboratories to CDRSS. Rapid influenza test data and respiratory syncytial virus data are acquired from facilities reporting rapid influenza tests via the National Respiratory and Enteric Virus Surveillance System (NREVSS) or CDRSS ILI module. Counts for cumulative totals begin with week ending October 8, 2016. Three week count data includes current week and two prior weeks. Data presented for RSV and rapid influenza testing represent information for the week prior to the current report week.
- 8. Daily visits and admissions associated with ILI from emergency department data is collected via EpiCenter and Hippocrates. Prior to these systems, data on ILI visits were only recorded one day per week usually on Tuesday. This system is maintained as a large amount of historical data allows for better seasonal comparisons.
- 9. Only LTCF outbreaks reported to NJDOH that receive an outbreak number are recorded in this report.
- 10. Data presented for New Jersey are for cases confirmed as of the current reporting week. Data presented for the United States represent data reported for the prior MMWR week. This data can be viewed at: <u>http://www.cdc.gov/flu/weekly/</u>.

	Influenza Activity	<u>Table 1</u> Level – Definitions for State A	Activity	
NJ Level	CSTE Level	<u> </u>	Definition	
		ILI Activity/Outbreaks		Lab Activity
	No Activity	ILI activity at or below baseline AND no detected outbreaks	AND	No lab confirmed cases
Low	Sporadic	Low ILI activity detected OR one lab confirmed outbreaks anywhere in the state	AND	Sporadic isolation of laboratory confirmed influenza
	Local	Increase in ILI activity OR two or more lab confirmed outbreaks in one public health region (Other regions not experiencing increased ILI activity)	AND	Recent (within 3 weeks) laboratory activity in the region with increased ILI
Moderate	Regional	Increase in ILI activity OR two or more lab confirmed outbreaks in at least 2 public health regions (Other regions not experiencing increased ILI activity)	AND	Recent (within 3 weeks) laboratory activity in the region with increased ILI
High	Widespread	Increase in ILI activity OR two or more lab confirmed outbreaks in > 2 public health regions	AND	Recent (within 3 weeks) laboratory activity in the region with increased ILI

	<u>Table 2</u> Influenza Activity Level – Definitions for Publ	ic Health	Regions
	<u> </u>	Definitior	<u>1</u>
<u>NJ Level</u>	ILI Activity/Outbreaks		Lab Activity
Low	Low ILI activity detected OR one lab confirmed outbreaks anywhere in the region	AND	Sporadic isolation of laboratory confirmed influenza anywhere in the region
Moderate	Increased ILI activity in less than half of the counties in the region OR two lab confirmed outbreaks in the public health region	AND	Recent (within 3 weeks) laboratory activity in same counties of the region with increased ILI
High	Increased ILI activity in more than half of the counties in the region OR three or more lab confirmed outbreaks in the region	AND	Recent (within 3 weeks) laboratory activity in more than half of the counties in the region with increased ILI

Notes:

ILI activity: Systems used to detect increases in ILI activity include: ILINet (i.e., sentinel providers), school absenteeism data, ED ILI visits and admissions collected via Hippocrates and EpiCenter systems, LTCF ILI data, LTCF outbreak data, and information on influenza mortality (122 city, influenza associated death report). Lab Activity: Virologic surveillance data from PHEL and commercial laboratories will be used as the primary data source for the above levels. However, rapid influenza test data will also be considered when determining the appropriate activity levels.

### INFLUENZA LABORATORY REPORTS BY COUNTY

### Counts represent total positive specimens from week ending October 7, 2016 to current MMWR week

### Source: CDRSS

Frequency			R	ESULT		
	COUNTY(COUNTY)	Influenza A - Typing not performed	Influenza A 2009 H1N1	Influenza AH3	Influenza B	Total
	ATLANTIC	464	0	32	33	529
	BERGEN	1316	1	219	138	1674
	BURLINGTON	283	0	8	31	322
	CAMDEN	725	0	32	32	789
	CAPE MAY	13	0	1	4	18
	CUMBERLAND	14	0	4	0	18
	ESSEX	342	1	54	36	433
	GLOUCESTER	200	0	42	9	251
	HUDSON	577	0	73	48	698
	HUNTERDON	104	0	26	10	140
	MERCER	217	0	19	14	250
	MIDDLESEX	295	0	39	32	366
	MONMOUTH	721	0	22	53	796
	MORRIS	405	0	81	38	524
	OCEAN	566	0	11	35	612
	PASSAIC	757	0	71	83	911
	SALEM	7	0	1	0	8
	SOMERSET	65	0	23	8	96
	SUSSEX	59	0	20	10	89
	UNION	399	1	112	47	559
	WARREN	97	0	48	14	159
	Total	7626	3	938	675	9242

### **INFLUENZA LABORATORY REPORTS BY REGION**

### Counts represent total positive specimens from week ending October 7, 2016 to current MMWR week

Frequency		Table of	REGION b	y RESULT		
			R	ESULT		
	REGION	Influenza A - Typing not performed	Influenza A 2009 H1N1	Influenza AH3	Influenza B	Total
	Central East	1981	1	184	167	2333
	Central West	386	0	68	32	486
	Northeast	2235	2	346	222	2805
	Northwest	1318	0	220	145	1683
	South	1706	0	120	109	1935
	Total	7626	3	938	675	9242

#### **Source: CDRSS**

The following is a breakdown of counties contained within each public health region: Northwest: Morris, Passaic, Sussex, Warren; Northeast: Bergen, Essex, Hudson Central west: Hunterdon, Mercer, Somerset Central East: Middlesex, Monmouth, Ocean, Union South: Atlantic, Burlington, Camden, Cape May, Salem, Cumberland, Gloucester

## NJ ACTIVE INFLUENZA-LIKE ILLNESS SURVEILLANCE STATISTICS SURVEILLANCE DATE: 02/21/2017





		Long Term Car	e		Schools		Hospi	al Emergency	Dept
COUNTY	# Enrolled	# Reports Rec'd	% ILI	# Enrolled	# Reports Rec'd	% Absent	# Enrolled	# Reports Rec'd	% ILI
February 21, 2017 MMWR WEEK	8								
ATLANTIC	6	0	0.00	43	26	6.09	4	4	4.19
BERGEN	10	4	0.12	37	5	6.30	6	6	4.04
BURLINGTON	8	4	0.00	95	50	5.53	4	3	4.75
CAMDEN	1	0	0.00	1	0	0.00	7	7	7.67
CAPE MAY	3	0	0.00	14	9	7.28	1	1	5.30
CUMBERLAND	5	4	1.04	12	10	9.22	3	3	3.54
ESSEX	7	4	0.00	4	0	0.00	8	7	4.77
GLOUCESTER	3	1	1.61	4	1	5.51	2	2	6.59
HUDSON	4	0	0.00	14	4	6.07	6	6	4.96
HUNTERDON	4	4	0.67	8	7	5.80	1	1	5.75
MERCER	3	0	0.00	22	7	3.21	5	4	8.60
MIDDLESEX	11	5	0.00	22	18	5.10	6	6	6.88
MONMOUTH	6	1	0.00	16	6	9.38	5	5	5.16
MORRIS	2	0	0.00	9	2	3.82	4	4	2.32
OCEAN	8	1	4.07	5	4	6.50	4	4	3.47
PASSAIC	10	1	0.00	29	2	6.58	3	3	4.85
SALEM	0	0	0.00	3	2	5.78	1	1	6.17
SOMERSET	4	0	0.00	23	13	4.80	1	1	7.02
SUSSEX	2	1	4.55	5	2	6.96	2	2	1.89
UNION	1	0	0.00	51	17	4.60	5	5	3.67
WARREN	5	3	0.00	20	11	5.18	2	2	5.63
NW Region	19	5	0.22	63	17	5.71	11	11	3.75
NE Region	21	8	0.08	55	9	6.17	20	19	4.60
CW Region	11	4	0.67	53	27	4.82	7	6	7.94
CE Region	26	7	0.43	94	45	5.34	20	20	5.10
South Region	26	9	0.82	172	98	6.24	22	21	5.91
State Total	103	33	0.43	437	196	5.77	80	77	5.20

# NJ ACTIVE INFLUENZA-LIKE ILLNESS SURVEILLANCE STATISTICS SURVEILLANCE DATE: 02/21/2017





	RSV	Tests	Rapid Flu Tests	
County	# Positive	Total Tests Performed	# Positive	Total Tests Performed
February 21, 2017 MMWR WEE	(8		T. T.	
ATLANTIC	1	17	69	267
BERGEN	3	16	186	587
BURLINGTON	0	0	5	42
CAMDEN	2	20	48	385
CAPE MAY	1	10	18	104
CUMBERLAND	2	10	0	0
ESSEX	10	185	125	734
GLOUCESTER	1	6	27	204
HUDSON	0	6	15	54
HUNTERDON	3	17	25	114
MERCER	1	3	87	313
MIDDLESEX	6	61	12	125
MONMOUTH	16	132	145	760
MORRIS	12	225	23	148
OCEAN	2	18	48	226
PASSAIC	6	39	77	271
SALEM	0	0	0	0
SOMERSET	0	0	0	0
SUSSEX	7	115	24	119
UNION	9	120	2	24
WARREN	6	86	26	142
NW Region	31	465	150	680
NE Region	13	207	326	1375
CW Region	4	20	112	427
CE Region	33	331	207	1135
South Region	7	63	167	1002
State Total	88	1086	962	4619